



Adult Vaccine Schedule

Hotline: 1-800-275-0659
www.immunize-utah.org

Vaccine ▼	Age ►	19-49 years	50-64 years	≥65 years
Tetanus, Diphtheria, Pertussis (Td/Tdap)		1 Td booster dose (every 10 years) Substitute 1 dose of Tdap for Td		
Human Papillomavirus (HPV)		3 doses (females 11-26 years)		
Measles, Mumps, Rubella (MMR)		1 or 2 doses (2 doses if born in 1957 or later)	1 dose	
Varicella (Chickenpox)		2 doses		
Zoster (Shingles)				1 dose
Influenza		1 dose (every year)	1 dose (every year)	
Pneumococcal (PPV)		1 – 2 doses (one-time only revaccination with at least 5 years between doses)		1 dose (5 years or more since last dose)
Hepatitis A		2 doses		
Hepatitis B		3 doses		
Meningococcal		1 or more doses (based on age, risk conditions and vaccine type)		

 Vaccines recommended for all adults in these age groups who have no evidence of immunity

 Vaccines recommended for adults with medical, job-related, lifestyle or other conditions that put them at higher risk

For vaccine details visit: www.immunize-utah.org/public/sched_adult.htm
Always discuss vaccines you wish to receive with a medical provider.

Adult Vaccine Recommendations for Medical and Other Conditions

Conditions ► Vaccine ▼		Weak immune system due to cancer, blood disorder, or cancer treatment with drugs or x-rays	Diabetes, heart disease, chronic lung disease, alcoholism	No spleen	Chronic liver disease, receiving clotting factor concentrates	Kidney disease or kidney failure, receiving blood dialysis	HIV infection	Health-care worker
Tetanus, Diphtheria, Pertussis (Td/Tdap)	X (Td only)	X	X	X	X	X	X	X
Human Papillomavirus (HPV)		X	X	X	X	X	X	X
Measles, Mumps, Rubella (MMR)			X	X	X	X	X	X
Varicella (Chickenpox)			X	X	X	X		X
Zoster (Shingles)			X	X	X	X		X
Influenza	X	X	X	X	X	X	X	X
Pneumococcal (PPV)	X	X	X	X	X	X	X	X
Hepatitis A	X	X	X	X	X	X	X	X
Hepatitis B	X	X	X	X	X	X	X	X
Meningococcal	X	X	X	X	X	X	X	X

X Vaccines recommended for adults with these conditions who have no evidence of immunity

X Vaccines recommended for adults with these conditions only if they have additional risk factors (medical, job-related, lifestyle or other conditions)

Vaccines should not be given to adults with these conditions

Use age groups and intervals on other side of card for recommended vaccines